

METHOD FOR ASSESSING THE RISK OF CARDIOVASCULAR DISEASE

FIELD OF THE INVENTION

5 The present invention is generally directed to a method for assessing the risk of cardiovascular disease (CVD), such as coronary heart disease (CHD), including myocardial infarction, and cerebrovascular disease in an individual, such as a human. Specifically the invention is directed to a method of identifying a condition in an individual in which condition an elevated serum or plasma high-density lipoprotein (HDL) concentration or HDL cholesterol concentration provides enhanced protection against cardiovascular disease. In addition, the invention provides a method of predictably treating an individual in order to enhance the plasma or serum HDL or HDL cholesterol of the said individual. Furthermore the invention provides a kit for carrying out the methods.

15 BACKGROUND OF THE INVENTION

A large number of prospective population studies have shown that elevation of high-density lipoproteins (HDL) is associated with a reduced incidence of coronary events, coronary mortality and atherosclerotic progression.^{1,2} The etiologic role of HDL in atherosclerosis and CHD has not, however, been confirmed in randomized clinical trials. The reasons why HDL elevating therapies do not consistently reduce cardiovascular risk are unknown.

There is a paradigm according to which any elevation of HDL is beneficial to health. This is, however, challenged by three lines of observations, which have been left unexplained. First, in populations with heavy alcohol intake, a high plasma HDL cholesterol concentration does not associate with reduced coronary and total mortality.³ Second, in alcoholics, a high HDL is not associated with effective reverse cholesterol transport.⁴ Third, recent reports suggest that a combination of a fibrate and cerivastatin, a HMG-CoA reductase inhibitor (Astatin®) might induce deaths, even though this combination

raises HDL levels. Common to both observations is that HDL elevation is caused by general liver induction or liver damage. Statins tend to elevate hepatic transaminases in plasma.⁵ Also alcohol elevates both HDL and liver transaminase levels. A wide variety of chemicals can produce liver enlargement, peroxisome proliferation, and induction of peroxisomal and microsomal fatty acid-oxidizing enzyme activities.

An undamaged liver has phase I and phase II detoxification systems. The phase I consists of the cytochrome P450 enzymes (CYP). Mutations in the genes that encode these enzymes reduce the efficacy of the CYP system and lead to the predisposition to liver damage. Also, gene mutations in the phase II detoxification enzymes lead to an enhanced sensitivity to liver damage. The phase II enzymes are defined here to include liver enzymes such as the catalase, paraoxonases, superoxide dismutases, glutathione peroxidases, glutathione synthases, glutathione reductases, glutathione transferases, glutamyl-cysteinyl synthase, quinone reductases, diaphorases, thioredoxins, glutaredoxins, peroxiredoxins, epoxide hydrolases, aldehyde hydrolases, aldo-keto reductases, properdins, selenoproteins P and W, N-acetyl-transferases, metallothioneins, sulfurtransferases, alcohol dehydrogenases, aldehyde dehydrogenases, glutamate dehydrogenases, dihydrodiol dehydrogenases, or carboxyl esterases. DNA mutations in any of the genes encoding these proteins can cause liver damage and impair the protective function of HDL.

Variation in the response to HDL elevating drugs may be due to genetic variations that may provide a molecular basis for differences in drug metabolizing enzymes such as CYP1, CYP2, and CYP3 subtypes.

Oxidative stress and free radicals have been implicated in the etiology of a number of diseases, including cancers, coronary heart diseases and type II diabetes. The human body has a number of endogenous free radicals scavenging systems, which have genetic variability. The serum paraoxonase (PON) is an enzyme carried in the HDL that contributes to the detoxification of organophosphorus compounds but also of carcinogenic

products of lipid peroxidation.⁶⁻¹⁴ *PON1* is polymorphic in human populations and different individuals also express widely different levels of this enzyme.^{9,11-13}

SUMMARY OF THE INVENTION

5

The object of the present invention is a method of identifying a condition in an individual in which an elevated serum or plasma HDL concentration or HDL cholesterol concentration provides enhanced protection against cardiovascular disease, the method comprising the step of testing the individual for a disorder that detrimentally affects the HDL function, i.e. the protective effect of HDL, whereby absence of such a disorder is an indication of enhanced protection against cardiovascular disease when said individual exhibits elevated serum or plasma HDL or HDL cholesterol concentration.

Furthermore, the invention is directed to a method of treatment of an individual to protect the individual against the risk of cardiovascular disease, the method comprising the steps of testing the said individual for a disorder which detrimentally affects the protective effect of HDL, identifying and selecting an individual free of said disorder, and treating the selected individual in order to enhance the HDL or HDL cholesterol level of said individual.

20

According to a further aspect the present invention provides a method for assessing the risk of cardiovascular disease in an individual, the method comprising the step of determining the serum or plasma HDL or HDL cholesterol concentration in said individual, and testing the individual for a disorder that detrimentally affects the HDL function, whereby identification of such a disorder is an indication of reduced protection against, i.e. an increased risk of cardiovascular disease when said individual exhibits elevated serum or plasma HDL or HDL cholesterol concentration.

In addition the invention is directed to a kit for use in the above methods, comprising means for testing the individual for a condition or disorder which affects the protective effect of HDL.

30

As is understood by the person skilled in the art, the HDL level in an individual can be assessed by determining the HDL concentration or a fraction thereof, e.g. the HDL cholesterol concentration of said individual.

5 DETAILED DESCRIPTION OF THE INVENTION

According to the invention, a condition or disorder which affects the protective effect or function of HDL, is, for example, liver damage or a condition involving oxidative stress. Both liver damage and oxidative stress have a detrimental effect on the protective effect
10 or function of HDL against cardiovascular disease, such as coronary heart disease, including myocardial infarction, and cerebrovascular disease.

The protective action of HDL depends on the ability of the liver to maintain the antioxidative capacity of HDL and the efficacy of HDL in the reverse transport of cholesterol
15 from the arteries to the liver. An elevation of γ -glutamyltransferase (GGT) indicates that these functions of HDL are compromised.

A condition of liver damage in an individual can be established in many ways, a convenient method involving determination of the serum or plasma activity or concentration of
20 an enzyme marker comprising γ -glutamyltransferase. In a condition involving liver damage, the concentration of γ -glutamyltransferase is elevated over the normal or reference values. This reference value or range can vary to some degree according to the specific methods used for determining the marker, but typically the reference value will be in the range of 20 to 100 units/L. For many purposes, a suitable value is 60 units/L. γ -
25 glutamyltranspeptidase (EC 2.3.2.2) acts as a glutathionase and catalyzes the transfer of the glutamyl moiety of glutathione to a variety of amino acids and dipeptide acceptors. This enzyme is located on the outer surface of the cell membrane. It is widely distributed in mammalian tissues involved in absorption and secretion. In humans, hepatic GGT activity is elevated in some liver diseases. GGT is released into the bloodstream
30 after liver damage.

Patients with cholestasis usually have increased serum γ -glutamyltransferase concentrations, and the concentrations may be increased by certain enzyme-inducing drugs or alcohol abuse. Measurements of serum γ -glutamyltransferase aid in interpreting elevated serum alkaline phosphatase values. γ -glutamyltransferase activity in serum is the sum of the activities of heterogeneous isoenzymes that migrate in zone electrophoresis as follows: GGT1 to the prealbumin-albumin region, GGT2 to the alpha-1-globulin region, GGT3 to the alpha-2-globulin region, and GGT4 to the beta-globulin region.

Instead of, or in addition to, measuring the γ -glutamyltransferase activity or concentration, it is possible to use genotyping of genomic DNA from a sample of said individual, and to identify mutations or polymorphisms in the DNA which influence liver damage or plasma or serum γ -glutamyltransferase activity or concentration. There are several γ -glutamyltransferase genes located on chromosome 22 and at least two of these appear to be transcribed. A third alternative is to measure the expression at the RNA level of the γ -glutamyltransferase.

The sample from which DNA can be extracted can be for example a blood sample. Genotyping can be carried out by using per se known techniques, for example PCR techniques involving the use of suitable primers and amplification systems. The genotyping method can be amplified restriction fragment length polymorphism (ARFLP) that utilizes PCR and restriction enzyme cleavage-site recognition. Additional methods such as DNA amplification by PCR followed by minisequencing and or sequence-specific oligonucleotide probe (SSOP) analysis can also be used. Also, genotyping can be performed by using DNA microarrays or DNA chips that provide information in the same assay of a number of DNA polymorphisms that affect the liver function. It is foreseen that a large number of DNA polymorphisms such as single nucleotide polymorphisms (SNP) are determined by the use of a single DNA chip. Also the expression of the genes encoding the γ -glutamyltransferase and the phase I and II detoxification enzymes can be assayed by microarray.

Oxidative stress is another condition which has a detrimental effect on the protective effect of HDL. A suitable marker for oxidative stress is the paraoxonase enzyme. The activity or concentration of paraoxonase can be determined in a serum sample from the individual, using per se known techniques, for example based on the capacity of paraoxonase to hydrolyse paraoxon, and by monitoring p-nitrophenol formation, for example using absorbance techniques. A reduced paraoxonase activity is an indication of oxidative stress, including increased lipid peroxidation. Consequently a low paraoxonase activity is an indication that the protective effect of HDL is impaired in the individual. A reference value within a reference range of 40 to 200 nmol/ml/min is usually applicable, a typical normal value for paraoxonase activity being appr. 100 nmol/ml/min.

Instead of, or in addition to, measuring the paraoxonase activity or concentration, it is possible to apply genotyping of DNA from a sample of said individual, and identification of mutations or polymorphisms which influence plasma or serum paraoxonase activity or concentration. Two polymorphisms are currently known in human *PON1*. The Q191R polymorphism was the first mutation of *PON1* reported.^{9,12} The second one is the missense mutation of A to T in codon 54, producing a substitution of methionine (M) to leucine (L) (Met54Leu⁸; known also as Met55Leu⁹). Both these polymorphisms have been shown to affect serum PON activity,^{12,15} and in particular, the L54 allele has been associated with an increased PON activity. A further alternative is to measure the expression of the genes encoding the PON enzyme.

According to the US patent 6,242,186, homozygosity of the L54 allele in the *PON1* gene protects against certain diseases associated with oxidative stress. The L allele has consistently been associated with an increased paraoxonase activity in human serum.⁷⁻¹² It was observed that there was less lipid peroxidation among men who carried the *PON1* 54 L allele. In such individuals, an enhanced HDL or HDL cholesterol concentration would therefore have a protective effect against cardiovascular disease. In the opposite, individuals who do not carry this mutation would not benefit from the protective effect of HDL against cardiovascular disease.

For genotyping purposes, DNA can be extracted for example from a blood sample. Genotyping can be carried out by using per se known techniques, for example PCR techniques involving the use of suitable primers and amplification systems. Such a system is described for example in the US 6,242,186.

The antioxidative capacity of HDL can be assessed by isolating HDL from plasma or serum e.g. by ultracentrifugation or precipitation and exposing the isolated HDL to oxidizing conditions e.g. by adding to the reaction oxidative agents such as oxygen free radicals such as peroxy radical, superoxide radical, hydroxyl radical or hydroperoxy radical. The radicals can be generated chemically utilizing the Fenton-Haberman-Weiss reaction for instance by adding reduced transition metal such as copper or iron, by using a radical generating substance such as ABAP (2,2'-azobis(amidinopropane) dihydrochloride) or AMVN (2,2'-azobis(2,4)-dimethylvaleronitrile) or by ionizing or other radiation, UV light, heating or by other means. The resistance of the target HDL (HDL isolated from an individual being examined) can be determined as the time lag to oxidation of HDL when exposed to said radicals. The oxidation of HDL can be determined by monitoring the formation of conjugated dienes at 234 nm absorbance by a spectrophotometer or by measuring periodically the concentration of an indicator compound of oxidation. Such a compound can be an oxidized phospholipid such as lysophosphatidylcholine (lysolecithine), an oxidized fatty acid such as hydroxy or epoxy fatty acid, or a cholesterol oxidation product such as hydroxy cholesterol or epoxy cholesterol or keto-cholesterol. The start of oxidation of HDL or the maximum rate of oxidation can be determined. The reference values are different for different methods. As an example, if oxidation of HDL is monitored spectrophotometrically following the formation of conjugated dienes at 234 nm, and copper ions are used to induce oxidation at a concentration of 10-100 micromoles per liter, a lag time of less than 30-200 min is an indication of reduced antioxidative capacity of HDL.

Lipid peroxidation in vivo can be assessed by measuring either immunologic response to immunogenic epitopes of oxidized lipoproteins, such as antibodies to oxidized low den-

sity lipoprotein.¹⁶ Lipid peroxidation in vivo can also be assessed by measuring oxidation products of lipids or lipoproteins such as oxidized phospholipids, oxidized fatty acids, or cholesterol oxidation products.¹⁶ Oxidized fatty acids such as hydroxy and epoxy fatty acids can be measured by gas chromatography mass spectrometry or immunochemical methods. Oxidation products of arachidonic acid such as isoprostanes can be used as indicators of lipid peroxidation in vivo. Lipid peroxidation can also be assessed by determining the proportion of electronegative LDL of total LDL by chromatographic or electrophoretic methods. Further, lipid peroxidation can be assessed by measuring plasma or serum concentration of conjugated dienes, an oxidation product of dienes. The reference values depend on the method used. As an example, plasma F₂-isoprostane levels of 20-60 ng/L or more, total plasma hydroxy fatty acids of 1-5 μ mol/L or more and plasma electronegative LDL of 3-10% or more of total LDL indicate increased lipid peroxidation in vivo.

The present invention also makes it possible to treat an individual in order to protect said individual against the risk of cardiovascular disease, by identifying whether said individual is responsive to the beneficial effects of a high HDL concentration. Such a method comprises a step of determining whether said individual has a condition which detrimentally affects the effect of high HDL. If said individual is free of such a condition, such individual can be treated in order to enhance his HDL level.

Such a treatment can be a drug treatment. A suitable drug can be a drug selected from the group consisting of niacin, a statin, an apolipoprotein AI or AII synthesis enhancing agent, a PPAR alpha agonist such as fibrate, a PPAR gamma or delta agonist, a sterol absorption inhibiting agent such as a resin, a CETP inhibitor, an ACAT inhibitor, a PLTP agonist, a LCAT agonist, a lipoprotein lipase (LPL) agonist, a hepatic lipase agonist, a scavenger receptor B1 (SRB1) agonist, or an ATP-binding cassette A1 (ABC1) agonist. A statin can be for example selected from the group consisting of atorvastatin, fluvastatin, lovastatin, pravastatin and simvastatin, a fibrate can be selected from the group consisting of bezafibrate, ciprofibrate, clofibrate, fenofibrate and gemfibrozil, and

a resin can be selected from the group consisting of colestipol and cholestyramin. It is, however, also possible to enhance HDL through physical activity or physical exercise.

5 The invention also provides for kits suitable for carrying out the methods according to the invention. Such a kit carries the necessary means for identifying a condition which affects the protective effect of HDL, such as for example the means necessary to determine enzyme, for example γ -glutamyltransferase or paraoxonase activity in a sample, such as a serum sample from the individual, or means for performing necessary genotyping of a DNA sample from said individual. In addition the kit can contain means
10 for measuring HDL or HDL cholesterol in a sample, such as a serum or plasma sample from the said individual. Such kits preferably contain the various components needed for carrying out the method packaged in separate containers and/or vials and including instructions for carrying out the method. Thus, for example, some or all of the various reagents and other ingredients needed for carrying out the determination, such as
15 buffers, primers, enzymes, control samples or standards etc can be packaged separately but provided for use in the same box. Instructions for carrying out the method can be included inside the box, as a separate insert, or as a label on the box and/or on the separate vials.

20 *Experimental*

In the following tests, the protective effect of HDL elevation in patients with liver damage was studied.

25 For assessing the protective effect, a prospective cohort study, the "Kuopio Ischaemic Heart Disease Risk Factor Study" (KIHD).^{1,2} was used. The study protocol for KIHD was approved by the Research Ethics Committee of the University of Kuopio, Finland. The study sample comprised men from Eastern Finland aged 42, 48, 54 or 60 years. A total of 2682 men were examined during 1984-89. All participants gave a written in-
30 formed consent. Relevant baseline measurements were available for 2464 men. The average follow-up time was 11.4 years resulting to over 28,000 person-years of follow-up.

γ -glutamyltransferase activity was determined according to the Nordic recommendation.¹⁷ The measurement of cholesterol concentration in serum lipoproteins and other risk factors, and the classification of acute coronary events and deaths have been described.^{1,2}

5

Among men whose liver enzyme (γ -glutamyltransferase) was within the normal range (60 IU/L or less), elevation of HDL was associated with decreased risk of acute coronary event (Table). On the average, the risk was reduced by 44% (95% confidence interval 14-68%) per each mmol/L of serum HDL cholesterol. However, in men whose
10 liver enzyme was elevated, the risk increased 3.3-fold (95% CI 1.2 to 9.3 -fold) per each mmol/L of HDL cholesterol. These relative risks differed significantly of each other ($p < 0.01$). The addition of any measured risk factor as a covariate singly or jointly did not affect this difference. Similarly, the relative risks for coronary, all cardiovascular and all-cause death were significantly different between men who had no liver enzyme
15 elevation and those who did (Table). There was a similar trend for cerebrovascular strokes.

As an indicator of lipid peroxidation, serum ferritin concentration was used. The study population was divided into those with normal serum ferritin (200 micrograms per liter or less) and those with elevated serum ferritin ($>200 \mu\text{g/L}$). A high serum HDL concentration was associated with a reduced cardiovascular mortality only in the subjects
20 whose serum ferritin was normal (relative risk 0.60, 95% CI 0.33 to 1.09, $p = 0.095$), whereas a high HDL tended to be associated with an increased risk (relative risk 1.02, 95% CI 0.37 to 2.77, $p = 0.976$) among those with elevated serum ferritin. There were
25 similar trends for the incidence of acute coronary events, cerebrovascular strokes and coronary deaths.

When the study cohort was stratified according to the PON1 codon 192 genotype, a high serum HDL cholesterol concentration was associated with a reduced risk of myocardial
30 infarction only among the wild type (arginine) homozygotes (relative risk 0.04, 96% CI

0.01 to 0.19, $p < 0.001$), whereas the associations of HDL cholesterol with myocardial infarction risk in subjects with the other genotypes were weak. There were similar trends in cerebrovascular strokes and cardiovascular and coronary deaths.

- 5 Our population-based data indicate that high serum HDL levels lose their protection against CHD among men who have liver damage, enhanced lipid peroxidation, a genotype that predisposes to liver damage or to enhanced lipid peroxidation. This effect modification was observed also for cardiovascular and total mortality, although high HDL was not protective of cerebrovascular strokes and cardiovascular and total mortality in our study. Our observations imply that an elevation of HDL is not always beneficial for human health. The liver damage and enhanced lipid peroxidation may be caused by heavy alcohol intake, drugs and hepatotoxic nutrients or contaminants in food.
- 10

Relative risk of acute coronary events, coronary, cardiovascular and any death, per 1 mmol/L of serum HDL cholesterol, in men without and with liver damage at baseline.

Outcome (number of men with each event)	No liver damage: γ -glutamyltransferase ≤ 60 IU/L or less (n=2253)			Liver damage: γ -glutamyltransferase >60 IU/L (n=211)			p for difference
	Relative risk	95% confidence interval	p-value	Relative risk	95% confidence interval	p-value	
Acute coronary event (n=381)	0.56	0.32, 0.86	0.008	3.32	1.19, 9.29	0.022	<0.01
Coronary death (n=141)	0.59	0.28, 1.24	0.161	5.16	1.23, 21.64	0.025	<0.01
Cardiovascular death (n=187)	0.91	0.49, 1.69	0.763	6.01	1.74, 20.80	0.005	<0.01
All-cause death (n=370)	0.95	0.61, 1.48	0.818	2.46	1.16, 5.22	0.019	<0.05

5 Cox' proportional hazards' models are adjusted for age, cigarette-years, serum apolipoprotein B (mg/L), use of antihypertensive drugs, maximal oxygen uptake (mL/kg x min), history of any atherosclerosis-related disease, family history of CHD and five examination years.

References

- 5 1. Salonen JT, Salonen R, Seppänen K, Rauramaa R, Tuomilehto J. High density lipoprotein, HDL₂ and HDL₃ subfractions and the risk of acute myocardial infarction: a prospective population study in Eastern Finnish men. - *Circulation* 1991; **84**: 129-39.
- 10 2. Salonen JT, Ylä-Herttuala S, Yamamoto R, Butler S, Korpela H, Salonen R, Nyyssönen K, Palinski W, Witztum JL. Autoantibody against oxidised LDL and progression of carotid atherosclerosis. *Lancet* 1992; **339**: 883-7.
- 15 3. Perova NV, Oganov RG, Williams DH, Irving SH, Abernathy JR, Deev AD, Shestov DB, Zhukovsky GS, Davis CE, Tyroler HA. Association of high-density-lipoprotein cholesterol with mortality and other risk factors for major chronic noncommunicable diseases in samples of US and Russian men. *Ann Epidemiol* 1995; **5**: 179-85.
- 20 4. Liinamaa MJ, Hannuksela ML, Kesäniemi YA, Savolainen MJ. Altered transfer of cholesteryl esters and phospholipids in plasma from alcohol abusers. *Arterioscler Thromb Vasc Biol* 1997; **17**: 2940-7.
- 25 5. Farmer JA, Torre-Amione G. Comparative tolerability of the HMG-CoA reductase inhibitors. *Drug Saf* 2000; **23**: 197-213.
- 30 6. Mackness MI, Thompson HM, Hardy AR, Walker CH. Distinction between 'A'-esterases and arylesterases. Implications for esterase classification. *Biochem J* 1987; **245**: 293-62
- 35 7. Mackness MI, Arrol S, Durrington PN. Paraoxonase prevents accumulation of lipoperoxides in low-density lipoprotein. *FEBS Lett* 1991; **286**: 152-4.
- 40 8. La Du BN, Adkins S, Kuo CL, Lipsig D. Studies on human serum paraoxonase/arylesterase. *Chem Biol Interact* 1993; **87**: 25-34
9. Humbert R, Adler DA, Disteché CM, Hassett C, Omiecinski CJ, Furlong CE. The molecular basis of the human serum paraoxonase activity polymorphism. *Nature Genet* 1993; **3**: 73-6.
10. Davies HG, Richter RJ, Keifer M, Broomfield CA, Sowalla J, Furlong CE. The effect of the human serum paraoxonase polymorphism is reversed with diazoxon, soman and sarin. *Nature Genet* 1996; **14**: 334.
11. Mackness MI, Mackness B, Durrington PN, Connelly PW, Hegele RA. Paraoxonase: biochemistry, genetics and relationship to plasma lipoproteins. *Curr Opin Lipidol* 1996; **7**: 69-76.
12. Mackness MI, Arrol S, Mackness B, Durrington PN. Alloenzymes of paraoxonase and effectiveness of high-density lipoproteins in protecting low-density lipoprotein against lipid peroxidation. *Lancet* 1997; **349**: 851-2.
13. Mackness B, Durrington PN, Mackness MI. Polymorphisms of paraoxonase genes and low-density lipoprotein peroxidation. *Lancet* 1999; **353**: 468-9.
14. Shih DM, Gu L, Xia Y-R, et al. Mice lacking serum paraoxonase are susceptible to organophosphate toxicity and atherosclerosis. *Nature* 1998; **394**: 284-7.

15. Garin MC, James RW, Dussoix P, et al. Paraoxonase polymorphism Met-Leu54 is associated with modified serum concentrations of the enzyme. A possible link between the paraoxonase gene and increased risk of cardiovascular disease in diabetes. *J Clin Invest* 1997; **99**: 62-6.
- 5 16. Salonen JT. Markers of oxidative damage and antioxidant protection: assessment of LDL oxidation. *Free Rad Res* 2000; **33** suppl.: S41-46.
- 10 17. The Committee on Enzymes of the Scandinavian Society for Clinical Chemistry and Clinical Physiology. Recommended method for the determination of gamma-glutamyltransferase in blood. *Scand J Clin Lab Invest* 1976; **36**: 119-125.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200